

# **Biophysical allergy therapy**

## **Retrospective study from a medical practise**

P. Schumacher  
Summary R. Moyses

### **Introduction**

Between September 1989 and October 1990, Dr. P. Schumacher of Innsbruck carried out a study of around 200 allergy patients which will be summarised here.

In his book *Biophysikalische Therapie der Allergien*<sup>1</sup> [Biophysical allergy therapy], in which this study is described in detail alongside numerous other useful tips on treating allergy, he writes: "Our own experience now covers well over 1000 such individual patients, all of whom have seen their allergy completely eliminated, no matter what form it took.

Added to these are the many patients treated by other colleagues. These colleagues - doctors and non-medical practitioners - learnt the method in our seminars and have been using it successfully in their practises, in some cases for several years now.

It is difficult to obtain a quantitative overview, yet the extensive feedback we receive shows that the method is also used elsewhere with considerable success. We felt it necessary, at least at the start of this development, to present some statistics independent of our personal experience."

B. Schneider<sup>2</sup> demonstrates that therapeutic action can also be verified through retrospective studies from medical practise when he suggests "assessing effectiveness by systematically observing success and failure in practise" as an alternative to conventional double blind studies. Schumacher published just such a study back in 1991, understandably causing a great sensation.

### **Patient selection**

All patients, mostly children and adolescents, undergoing *BICOM* allergy therapy within a 6-month period were recorded on a list, regardless of age, diagnosis, severity of the disease and type of allergen. To be included in the study diagnosis of allergy had to be confirmed by one or more allergy tests and there had to be an identifiable and confirmed link between the allergy and allergic reaction. So the study group was not specifically selected but consisted of the spontaneous patient body within a certain period of time.

Scientific studies

Copyright C 1999 Institut für Regulative Medizin, D-82166 Gräfelfing

A total of 164 patients with 204 cases of treatment were recorded. Results were gathered using a questionnaire. Questionnaires were sent out 5 months after treatment was completed. Patients with several allergies received a separate questionnaire for each allergen treated so that each series of treatments could be assessed separately.

The investigation lasted 6 months so that the interval between treatment and feedback was 5 months minimum and, at most, 11 months. 200 of the 204 questionnaires sent out were returned, giving a 98% response rate.


Patients were asked about tolerance, side-effects of treatment, relapses and overall assessment of the success of treatment. Since the study mainly involved children and adolescents, most of the questionnaires were completed by parents.

The tables indicate age distribution and frequency with allergies were detected. As expected, wheat and cows' milk were the most common food allergies with goose down in the case of inhalation allergens. Hay fever was excluded from the program due to its special peculiarities.

<sup>1</sup> Sonntag Verlag, Stuttgart, 1994;

<sup>2</sup> Prof. B. Schneider, Hanover Medical School: Erfahrungsheilkunde 11/1994;


### age distribution - 200 cases



## Treatment

Allergy treatment took the same form for all patients, namely just one *BICOM* therapy with the inverse oscillations of the allergen diagnosed and 8-fold amplification. Complete allergen abstinence was a pre-condition of starting treatment.

Abstinence could be stopped once treatment was completed. After treatment, the patient no longer had to avoid contact with the allergen but was instructed to report any further allergic reaction immediately.



## Evaluation of therapeutic results

The patients or their families carried out their own evaluation. They could choose from the following criteria:

1. The allergy has disappeared, i.e. since completing treatment the patient has tolerated the allergen without

Scientific studies

Copyright C 1999 Institut für Regulative Medizin, D-82166 Gräfelfing

suffering a reaction. The allergic reactions observed previously have not reappeared despite contact with the allergen.

2. The allergy has improved, i.e. symptoms of the allergy are still discernible but gradually much less marked.

3. The allergy continues unchanged. There is no discernible sign of success.

4. There has been a relapse, i.e. although the symptoms initially disappeared, the allergy to the same allergen has recurred.

5. It is not possible to assess the therapeutic result at the moment.


## Results

83.0% of cases were treated successfully, i.e. the reactions observed previously no longer occurred despite contact with the allergen. The patient was able to tolerate the allergen permanently and unrestrictedly without a reaction.

11.0% of cases had improved. The symptoms of the allergy were still discernible but gradually much less marked.

4.5% of cases were therapeutic failures. There was no discernible sign of success or the same allergy recurred despite initial improvement.

1.5% of cases could not be assessed.



## **Discussion**

With younger patients who display a better reactive state and who do not have additional severe stresses, preparatory, supporting or additional therapeutic measures are not necessary. A different therapeutic approach is often required for adults with multiple stresses.

The group assessed as "improved" comprised a total of 22 cases. Closer examination revealed that the symptoms still present had been misinterpreted and were only indirectly associated with the actual allergy process. For example, 8 cases in this group were patients with neurodermatitis who still displayed residual mycoses in the cutaneous areas affected, yet the allergy-induced neurodermatitis which had been the original cause was healed.

The follow-up check on the remaining 14 patients in this group revealed that either non-specific symptoms had been confused with allergy-induced symptoms, such as non-specific coughing due to infection in asthma patients or non-specific abdominal pain following completely cured allergic colitis, or that symptoms were still present from allergies which had not been identified.

A follow-up check of the group of therapeutic failures with no change in symptoms or with relapses revealed that, in 7 cases, abstinence had lapsed during the treatment phase, i.e. the patient was not kept strictly and consistently apart from the allergen. In 2 cases in this group there was evidence of a continuous lapse.

# Two years' experience with meridian-based allergy therapy Retrospective study from a medical practice

J. Hennecke

## 1. Introduction

Originally, an allergy could only be treated effectively by deactivating the allergy process through abstaining from the allergen for an indeterminate period of time. For purely technical reasons, about six treatment sessions were then also required. Therapeutic failures arose if the patient could not observe the abstinence period strictly enough or allergy abstinence was not possible at all.

The idea of treating the allergy via disturbed acupuncture meridians originates from kinesiology. I transferred this idea to *BICOM* therapy. This approach does not use physically generated inverse oscillations, instead energy blocks are released by stimulating certain acupuncture points.

Stimulabon was carried out with the *BICOM* device with the pathological information being picked up from an immunocompetent region of the body (cervical spine area) and directed to the appropriate points after being modulated through a H+Di circuit (program 530). Allergen abstinence was not necessary with this type of treatment and one therapy session was generally sufficient.

With considerable creativity a number of therapy versions were developed, all operating, however, according to the same basic principle: "energetic hyposensitisation" by removing energy blocks while in contact with allergens. However, experience soon showed that "energetic" therapy is easily disrupted by other energetic blocks, such as geopathy, scar interference fields, emotional blocks leading to a recurrence of the allergy treated previously.

For this reason, a therapeutic approach was developed whereby, using a series of connected *BICOM* programs, the patient would be placed energetically in a position to start allergy therapy and achieve effective and lasting release from allergy.

A pilot study was carried out on patients in my practice to objectivise therapeutic results and uncover and improve on the approach's weaknesses.

## 2. Method

The study comprised patients (mainly adults) who had been treated with *meridian-based allergy therapy* in my practice between June 1991 and 1993 and whose course of treatment had been completed.

### Allergy test

The allergens were determined with a kinesiological muscle test (generally *Deltoidaeus medialis*). Only allergens detected by this method were treated, regardless of any previous skin or RAST test. Substances from *Dr. Schumacher's* test sets were tested, together with substances brought by the patients and some of my own

Scientific studies

Copyright C 1999 Institut für Regulative Medizin, D-82166 Gräfelfing

ingredients (e.g. vaccines). With small children the test was conducted through a surrogate. Following therapy, the patient was also tested kinesiologically on all allergy contact zones (abdomen, costal arch, thymus, temples).

## **Allergy treatment**

*Meridian-based allergy therapy* was only carried out according to the original recommendation and was generally delegated to staff.

Device input: flexible electrode cervical spine area.

Device output: two button electrodes to stimulate the start and end point of the bladder, kidney, stomach and spleen-pancreas meridian, each for about one minute.

Allergen: on contact zone beneath the navel (conception vessel 6), if repeating treatment, also thymus or temples.

The other versions of allergy therapy were not routinely conducted so as not to make it harder to evaluate the therapeutic results.

As many pre-tested allergens as possible were treated, regardless of whether central allergens (milk, wheat), symptom-triggering allergens (house dust, pollen) or allergic stress (vaccination, pesticides, heavy metals).

## **Additional programs**

The treated allergies reappeared after a while if a geopathic stress or a scar interference field was ignored. These obstacles to treatment were always tested out kinesiologically and treated before each session of allergy therapy. The patient was also naturally informed about the need to cleanse the area where they sleep of harmful effects. The *geopathy balancing* and *elimination of scar interference* programs were used for this.

All patients were tested kinesiologically whether they accepted treatment from a psychological viewpoint, i.e. whether they wanted to be healthy. Where "psychological reversal" of this kind was present with the result that the patient was not prepared to seek to be healthy, a special program (991) was conducted to prepare for allergy therapy with rescue drops in the input cup and the patient connected only to the device output.

To stabilise general metabolism, a non-specific metabolic program *brief therapy for nutrient points* was also conducted before each allergy treatment session to compensate for impaired processing in the vitamin and mineral balance.

## **Additional treatment**

**Intestinal mycoses:** most allergy sufferers exhibit intestinal colonisation with *Candida* fungi and the consequential damaging effects on the immune system associated with the intestinal wall.

In addition to an antifungal diet, a three- to four-week course of nystatin (after prior therapy for lactose allergy)

Scientific studies

Copyright C 1999 Institut für Regulative Medizin, D-82166 Gräfelfing

was prescribed and the intestinal flora were subsequently built up (symbioflor, mutaflor or omniflora).

**Amalgam stress:** all patients were alerted to a possible link between toxic stress from mercury and the occurrence of allergies through damage to the immune system. For financial reasons, only a few patients were able to decide on amalgam cleansing. If amalgam cleansing was carried out, this was followed by *BICOM* elimination therapy and a prolonged course of selenium, zinc, vitamin C and homeopathic remedies.


### **3. Evaluation of the study**

Patients experienced some difficulty in assessing the success of their therapy and completing the questionnaire. In the case of symptom-triggering allergens, the patient can check the therapeutic result immediately. With masked allergies and allergic stress factors where therapy does not involve abstinence, the connection between allergen and symptoms is not obvious to a patient. Consequently he cannot judge whether he is insensitive to a single allergen; he can only judge whether, following completed treatment, he is symptom-free, his condition is improved or unchanged. The situation can only be clarified by our own testing and checking.

A questionnaire was used which was sent to each patient either immediately after completing a course of treatment or in May 1993. The first 200 questionnaires completed and submitted were analysed. Fig. 1 shows the frequency distribution of the allergens treated.

### **4. Experience of treating various allergens**

A report published for *BICOM* users contains detailed information on the peculiarities of treating allergies induced by house dust, animal epithelial cells, foodstuffs, contact, medication, etc<sup>1</sup>.



Tab.1: Allergen frequency – Incidence of allergens treated in 200 cases

<sup>1</sup> J. Hennecke: Zwei Jahre Erfahrungen mit der Allergie-Therapie ohne Karenz [Two years' experience of allergy therapy without abstinence], RTI volume 13 (papers, therapy tips, information), Institut für Regulative Medizin

## 5. Results

Since treating the individual syndromes achieved different results, it was important and interesting to record these separately (table 2). The overall result can be seen in figure 1.

**Allergically conditioned skin diseases** - This group contains neurodermatitis, endogenous eczema and other forms of chronic eczema, between which even dermatologists experience difficulty distinguishing. Yet this does not matter for the diagnosis and treatment of the underlying masked central allergy. We always encounter one or more central allergies such as milk, wheat or Candida, an indeterminate number of superficial foodstuff allergens and often also symptom-aggravating contact allergens such as tap water, house dust, pollen, clothing, cleaning products and ointments.

The longer the disease has been in existence and the older the patient, the more complex the therapy as the number of allergens usually increases, frequently changes and also non allergy-induced components are more common. Added to which, stresses from amalgam, toxins, viruses and superinfection of the cutaneous areas by bacteria and fungi complicate the picture. Unfortunately the results from treating adults with neurodermatitis are not as good as those for most children.

	Total	No change		Improved		Symptom-free	
Allergic skin diseases	68	8	12%	21	30%	39	58%
Pruritis	20	3	15%	10	50%	7	35%
Allergic conjunctivitis	16	1	6.5%	5	31%	10	62.5%
Allergic intestinal disorders	13	2	15.5%	1	7.5%	10	77%
Allergic respiratory tract diseases	46	6	13%	21	45.5%	19	41.5%
Pollen allergy	69	16	23%	21	31%	32	47%
Total		15.5%		34.1%		50.4%	

**Tab. 2:** Result broken down according to individual syndromes

If treatment is carried out without abstinence, it is almost impossible for the patient to evaluate the immediate success as there is no unmasking or deterioration in their condition on contact with the allergen. Ultimately they can only judge whether their cutaneous manifestations heal completely without being able to attribute this to the treatment of a specific allergen. This fact obviously makes it harder to evaluate the completed questionnaires. It is only in classic contact eczema that the connection with the triggering allergen is evident.

Similar mechanisms to those described above underlie the various types of allergically induced exanthema and we have therefore included them here. Treatment is usually less problematical and easier to evaluate, if the triggering allergens are known or were found.

Scientific studies

Copyright C 1999 Institut für Regulative Medizin, D-82166 Gräfelfing

Of the 200 cases investigated, 68 patients had cutaneous symptoms. Of these, 39 were symptom-free (58%), 21 improved (30%) and 8 showed no change (12%) following therapy (table 2).

### **Pruritis**

Masked allergies play a major part in itching with no visible rash, but other disorders such as diabetes, liver and kidney disorders and psychological factors also cause this condition. An allergen which is found and treated is often just one aspect of the problem. Of 20 patients with pruritis, 7 were symptom-free (35%), 10 improved (50%) and 3 showed no change (15%) (table 2).

### **Allergic conjunctivitis (all year)**

Allergically induced conjunctivitis may occur in isolation or linked with other allergic symptoms. It quite frequently accompanies palpebral eczema (often wheat allergy) and allergic sinusitis. Of 16 patients with allergic conjunctivitis (none of whom were allergic to pollen), 10 were symptom-free (62.5%), 5 improved (31%) and 1 showed no change (6.5%) (table 2).

### **Allergic intestinal disorders**

This group includes chronic diarrhoea, chronic constipation, non-specific colitis, irritable bowel syndrome, Roemheld's syndrome, umbilical colitis and so-called "functional abdominal pain." The intestine is almost always colonised by Candida and must be treated in addition to allergy therapy. Allergies to milk, lactose, wheat, yeast and Candida play a major part. Foodstuff allergies are always present in Colitis ulcerosa and Crohn's disease. Allergy therapy alone is not sufficient to treat the problem, however. Of 13 patients, 10 were symptom-free (77%), 1 improved (7.5%) and 2 showed no change (15.5%).

### **Allergic respiratory tract diseases (all year)**

This group includes bronchial asthma, chronic obstructive bronchitis, recurrent spastic bronchitis, chronic cough as well as chronic sinusitis. Here too, milk and wheat represent the most common central allergens. Frequently encountered symptom-inducing inhalation allergens are house dust, mould, down, feathers, animal hair as well as dust and solutions to which the individual is exposed at work. Foods (apples, nuts) also trigger shortness of breath. The patient can generally easily fudge whether they no longer react to these allergens or whether the reaction has diminished, although masking occurs here too. Factors unrelated to allergy (physical and emotional stress, fungal infection) play a large part with this group of patients. Of 46 patients, 19 were symptom-free following therapy (41.5%), 21 improved (45.5%) and 6 showed no change.

### **Pollen allergy**

In our practice, pollen allergy sufferers represented the largest and unfortunately the most difficult patient group. We waited at least one pollen season before analysing the results. Following therapy, 32 of the 68 patients were

Scientific studies

Copyright C 1999 Institut für Regulative Medizin, D-82166 Gräfelfing

symptom-free (47%), 21 improved (31%) and 16 showed no change (23%). There are a number of important reasons for this not wholly satisfactory result which are important for *BICOM* users and which are dealt with in a separate report<sup>2</sup>.

### **Other indications**

As well as the classic allergic disorders, a range of syndromes exist in which allergens also play a part, contrary to educated opinion. These include migraine, rheumatic disorders, psoriasis and colitis. Allergy treatment can only represent a corner-stone of the total therapy however. Yet cases arose where allergy therapy alone cured the condition. Migraine which had been present for a year disappeared completely with one treatment session for lactalbumin allergy. In other cases, a food allergy was the cause of impaired vision, arthritis, back pain, cardiac dysrhythmia, finger stiffness, tenderness, dizziness and depression.

<sup>2</sup> J. Hennecke: Zwei Jahre Erfahrungen mit der Allergie-Therapie ohne Karenz [Two years' experience of allergy therapy without abstinence], RTI volume 13 (papers, therapy tips, information), Institut für Regulative Medizin:

### **6. Side effects**


Of 200 patients, 6 reported extreme tiredness and physical exhaustion following therapy. 4 reported an immediate improvement in their overall condition. 6 patients experienced a deterioration in their allergy symptoms in the form of an initial exacerbation. Rashes intensified in 4 patients and 2 patients experienced increased coughing. All adverse side-effects disappeared within a few days.

### **7. Summary and discussion**

The overall therapeutic result of our pilot study on 200 allergy sufferers is as follows:

- Symptom-free
- Improved
- No change

### Therapeutic results of 200 cases treated



Considering these were mainly patients who had been treated extensively with conventional medicine and were still suffering from the side-effects of this, these results are impressive. However, it is these very problem patients who we, as BICOM therapists, want to endeavour to treat in a more satisfactory fashion.

Unsatisfactory therapeutic results and therapeutic failures may be caused by following:

*1. Hypersensitivity to the allergen has not completely disappeared*

In certain circumstances, treatment must be carried out twice or three times in the case of very strong allergens. If this is still unsuccessful, a different version of therapy should be tried. Therapy blocks should also be looked for.

*2. Hypersensitivity to the allergen disappears at first, but then recurs*

Here too therapy blocks are generally present (geopathy, scar interference fields, emotional blocks).

*3. Hypersensitivity to the allergen disappears, persistent or recurring symptoms are caused by another allergen*

The only solution here is to search thoroughly for new allergens and test and check them.

*4. Hypersensitivity to the allergen is eliminated and the symptoms are largely caused by other factors*

It is primarily a case here of toxin stress, mould infestation and intestinal dysbiosis. Our main task in the next few years will be to focus on this area.

Scientific studies

Copyright C 1999 Institut für Regulative Medizin, D-82166 Gräfelfing

Treatment of people with few allergies and small children generally produces better results than with adults suffering from multiple allergies. The more the therapy can be tailored to the individual, the better the results. Smaller practices with intensive patient care have the advantage here. Busy practices can only cope with the large and ever increasing number of allergy patients using methods which can be applied routinely and delegated.

Many good approaches can definitely be improved further. All *BICOM* therapists are also amateur researchers and, only by all users actively exchanging their experiences, will we find ways in future to satisfy all our allergy patients.

## **Clinical study on the use of BICOM resonance therapy on overstrain syndrome in top sportsmen**

Dr. med. Breda Jesensek Papcz, Dr. med. prim. Joze Barpvic  
German translation and adaptation T. Rome and R. Moyses

### **Introduction**

The organism's biophysical processes are fundamental characteristics which are present in all organisms and are superordinate to biochemical processes. The frequency range extends from very short to the extremely long wave range. The body creates a dynamic equilibrium between physiological and pathological oscillations through auto-regulation. A disturbed auto-regulation system manifests itself as disease.

The principle of *BICOM* resonance therapy consists of acting on processes such as this with endogenous physiological and inverted pathological electromagnetic oscillations in order to restore the body's dynamic equilibrium.

The purpose of this study is to test the effectiveness of *BICOM* resonance therapy on overstrain syndrome in top sportsmen compared with the conventional treatment.

### **Research group**

The study was carried out by Dr. Med. Brads Jesensek Papcz and Dr. med. prim. Joze Barocic under the direction of Prim. Dr. Sci. Dr. Med. Zmago Turk at the Maribor teaching hospital in Slovenia. The sportsmen with overstrain syndrome (footballers, athletes, sprinters, hurdlers, etc.) were divided into two groups, each containing 12 individuals.

Control group 1 was treated with conventional methods such as ultrasound on the pain site, electro-stimulation, cryotherapy and locally also with antirheumatic agents (voltaren gel).

Experimental group 2 was treated with *BICOM* resonance therapy, namely basic program and *sports injury* follow-up program with the therapy parameters adapted to the individual in accordance with energetic testing. Individually treated bioresonance oil was also applied locally.

Both groups of sportsmen were recommended to take a break from training to relieve the body until the syndrome was alleviated.

### **Assessing pain and therapeutic results**

Scientific studies

Copyright C 1999 Institut für Regulative Medizin, D-82166 Gräfelfing

The success of the treatment was evaluated using a visual analogue scale (VAS) and accompanying clinical status. The VAS is a simple, one-dimensional scale commonly used in medicine. To determine the severity of the pain, it is divided into 10 stages from 0 (no pain) to 10 (maximum conceivable pain). Admittedly the pain is perceived subjectively but this subjective estimation is necessary to judge how successful therapy has been and has proved useful in practice. It is simple and quick and consequently the most widely used method for determining the severity of pain and how treatment is progressing.

To carry out an accurate assessment, it is first carefully explained to the patient that stage 0 means "no pain" and that 10 is the maximum pain rating the patient can imagine for his situation.

## **Treatment**

**Control group 1:** this group was treated as follows:

1. Ultrasound treatment - 1 MHz/intensity 0.65 W/cm<sup>2</sup> for 5 to 10 mins, 5 times per week
2. Cryotherapy was carried out locally for up to 2 mins with the Kryojet device
3. Electro-stimulation - according to the customary standards - CP modulation (direct current) 5 mins and LP modulation (low frequency) 5 mins
4. The sportsmen also carried out individual cryotherapy at home according to the physiotherapist's directions
5. Antirheumatic agents (voltaren gel) were also used locally.

**Experimental group 2:** with this group, *BICOM* resonance therapy, type 4.0 device, was carried out as follows:

1. An introductory *basic program*

The basic program was selected in accordance with the conductivity tested

2. Sports *injuries* as follow-up program

Therapy type H + Di/bandpass setting 114kHz/amplification H 2.5 fold - Di 26 fold wobbling bandpass - interval operation - therapy time 5 min

The amplifications for H and Di were tailored to the individual patient by testing beforehand. The preset 5-minute therapy time was increased to 10 minutes for all patients and then adjusted more precisely. The parameters were not retested for each subsequent therapy session but those tested at the start were used.

## **Results**

Scientific studies

Copyright © 1999 Institut für Regulative Medizin, D-82166 Gräfelfing

The following tables indicate the relative success of the treatments used:

	Average VAS score prior to treatment	Average VAS score after treatment
Control group 1	5.25 (min 3/max 8)	2.6 (min 0/max 6)
<i>BICOM</i> group 2	5.41 (min 3/max 8)	0.61 (min 0/max 3)

**Tab.1:** Average pain rating

	Total treatment period in days	Number of treatment sessions
Control group 1	144	120
<i>BICOM</i> group 2	104	48

**Tab. 2:** Total treatment period and number of treatment sessions

There is a statistically significant difference between the groups of  $p < 0.05$ . All the parameters measured favour *BICOM* resonance therapy.

## Discussion

The results are surprising and favour *BICOM* resonance therapy. Better results were achieved in less time and with fewer sessions than using standard methods. As a result, it was not necessary to interrupt training for so long.

Within 12 months of their initial injury, 4 of the sportsmen in the *BICOM* group appeared for further treatment due to a recurrence. This time their condition was relieved after just two *BICOM* sessions.

The test group only included active top sportsmen who sustain injuries in the form of overstrain syndrome relatively frequently and who had repeatedly undergone several courses of physiotherapy prior to *BICOM* therapy. Despite this, their subjective assessment of *BICOM* therapy was positive.

## Summary

In conclusion, we can confirm that *BICOM* resonance therapy can be used successfully on overstrain syndrome without any difficult or negative side-effects.